

# New use of global warming potentials to compare cumulative and short-lived climate pollutants

Myles R. Allen<sup>1,2\*</sup>, Jan S. Fuglestedt<sup>3</sup>, Keith P. Shine<sup>4</sup>, Andy Reisinger<sup>5</sup>, Raymond T. Pierrehumbert<sup>2</sup> and Piers M. Forster<sup>6</sup>

**Parties to the United Nations Framework Convention on Climate Change (UNFCCC) have requested guidance on common greenhouse gas metrics in accounting for Nationally determined contributions (NDCs) to emission reductions<sup>1</sup>. Metric choice can affect the relative emphasis placed on reductions of 'cumulative climate pollutants' such as carbon dioxide versus 'short-lived climate pollutants' (SLCPs), including methane and black carbon<sup>2–6</sup>. Here we show that the widely used 100-year global warming potential (GWP<sub>100</sub>) effectively measures the relative impact of both cumulative pollutants and SLCPs on realized warming 20–40 years after the time of emission. If the overall goal of climate policy is to limit peak warming, GWP<sub>100</sub> therefore overstates the importance of current SLCP emissions unless stringent and immediate reductions of all climate pollutants result in temperatures nearing their peak soon after mid-century<sup>7–10</sup>, which may be necessary to limit warming to "well below 2 °C" (ref. 1). The GWP<sub>100</sub> can be used to approximately equate a one-off pulse emission of a cumulative pollutant and an indefinitely sustained change in the rate of emission of an SLCP<sup>11–13</sup>. The climate implications of traditional CO<sub>2</sub>-equivalent targets are ambiguous unless contributions from cumulative pollutants and SLCPs are specified separately.**

Establishing policy priorities and market-based emission reduction mechanisms involving different climate forcing agents all require some way of measuring what one forcing agent is 'worth' relative to another. The GWP<sub>100</sub> metric has been widely used for this purpose for over 20 years, notably within the UNFCCC and its Kyoto Protocol. It represents the time-integrated climate forcing (perturbation to the Earth's balance between incoming and outgoing energy) due to a one-off pulse emission of one tonne of a greenhouse gas (GHG) over the 100 years following its emission, relative to the corresponding impact of a one-tonne pulse emission of CO<sub>2</sub>. The notion of a temporary emission pulse is itself a rather artificial construct: it could also be interpreted as the impact of a delay in reducing the rate of emission of a GHG (see Methods).


This focus on climate forcing and a 100-year time horizon in GWP<sub>100</sub> has no particular justification either for climate impacts or for the policy goals of the UNFCCC, which focus on limiting peak warming, independent of timescale. Although it could be argued that, given current rates of warming, the goal of the Paris Agreement<sup>1</sup> to limit warming to 'well below 2 °C' focuses attention on mitigation outcomes over the next few decades, this focus is only implicit and presupposes that this goal will actually be met. Individual countries may also have goals to limit climate impacts

in the shorter term. These are acknowledged by the UNFCCC, but not quantified in terms of, for example, a target maximum warming rate. Metric choice is particularly important when comparing CO<sub>2</sub> emissions with SLCPs such as methane and black carbon aerosols. Black carbon has only recently been introduced into a few intended NDCs<sup>14</sup>, but may become increasingly prominent, as some early estimates<sup>15</sup> assign it a very high GWP<sub>100</sub>, even though the net climatic impact of processes that generate black carbon emissions remains uncertain<sup>16</sup> and policy interventions to reduce black carbon emissions are likely to impact<sup>6</sup> other forms of pollution as well. Here we combine the climatic impact of black carbon with that of reflective organic aerosols using forcing estimates from ref. 16 (see Methods).

At least one party to the UNFCCC has argued<sup>17</sup> that using the alternative global temperature-change potential (GTP) metric would be more consistent with the UNFCCC goal of limiting future warming. In its most widely used 'pulse' variant<sup>2</sup>, the GTP represents the impact of the emission of one tonne of a GHG on global average surface temperatures at a specified point in time after emission<sup>18</sup>, again relative to the corresponding impact of the emission of one tonne of CO<sub>2</sub>. Figure 1 shows how both GTP and GWP values for SLCPs such as methane and black carbon depend strongly on the time horizon. For long time horizons, SLCP GTP values also depend on the response time of the climate system, which is uncertain<sup>19,20</sup>. This latter uncertainty is a real feature of the climate response that is not captured by GWP, and so is not itself a reason to choose GWP over GTP. Other metrics and designs of multi-gas policies have been proposed<sup>21,22</sup>, some of which can be shown to be approximately equivalent to GWP or GTP<sup>23</sup>, but because only GWP and GTP have been discussed in the context of the UNFCCC, we focus on these here.

For any time horizon longer than ten years, values of the GTP are lower than corresponding values of the GWP for SLCPs. The time horizon has, however, a different meaning between the two metrics: for GWP it represents the time over which climate forcing is integrated, whereas for GTP it represents a future point in time at which temperature change is measured. Hence, there is no particular reason to compare GWP and GTP values for the same time horizon. Indeed, Fig. 1 shows that the value of GWP<sub>100</sub> is equal to the GTP with a time horizon of about 40 years in the case of methane, and 20–30 years in the case of black carbon, given the climate system response times used in ref. 16, for reasons given in the Methods<sup>24</sup>. Values of GWP and GTP for cumulative pollutants such as nitrous oxide (N<sub>2</sub>O) or sulfur hexafluoride (SF<sub>6</sub>) are determined primarily

<sup>1</sup>Environmental Change Institute, School of Geography and the Environment, University of Oxford, South Parks Road, Oxford OX1 3QY, UK. <sup>2</sup>Department of Physics, University of Oxford, Parks Road, Oxford OX1 3PU, UK. <sup>3</sup>Center for International Climate and Environmental Research - Oslo (CICERO), PO Box 1129 Blindern, 0318 Oslo, Norway. <sup>4</sup>Department of Meteorology, University of Reading, Earley Gate, PO Box 243, Reading RG6 6BB, UK. <sup>5</sup>New Zealand Agricultural Greenhouse Gas Research Centre, Private Bag 11008, Palmerston North 4442, New Zealand. <sup>6</sup>School of Earth and Environment, Maths/Earth and Environment Building, The University of Leeds, Leeds LS2 9JT, UK. \*e-mail: myles.allen@ouce.ox.ac.uk



**Figure 1 | GWP and GTP as a function of the time horizon.** **a**, Values for methane. **b**, Values for combined organic and black carbon. Solid lines show metrics calculated using current IPCC response functions<sup>16</sup>; dotted blue lines show the impact of varying the climate response time (see Methods). Dotted black lines show the value of  $GWP_{100}$ .

by forcing efficiencies, not lifetimes, and are hence similar to each other and almost constant over all these time horizons<sup>16</sup>. So, for a wide range of both cumulative and short-lived climate pollutants,  $GWP_{100}$  is very roughly equivalent to  $GTP_{20-40}$  when applied to an emission pulse, making it an approximate indicator of the relative impact of a one-off pulse emission of a tonne of GHG or other climate forcing agent on global temperatures 20–40 years after emission. The inclusion of feedbacks between warming and the carbon cycle can substantially increase GTP (and also, to a lesser degree, GWP) values, particularly on century timescales<sup>25</sup>. Here we follow the traditional approach, used for the most widely quoted metric values in ref. 16, of including these feedbacks in modelling  $CO_2$  but not other gases.

Figure 2a shows the impact on global average temperature of a pulse emission of various climate pollutants, with the size of the pulse of each gas being ‘equivalent’ (in terms of  $GWP_{100}$ ) to total anthropogenic  $CO_2$  emissions in 2011 (38 Gt $CO_2$ ): hence the pulse size is  $38/GWP_{100}$  billion tonnes of each forcing agent. SLCPs with high radiative efficiencies, such as methane, black carbon and some hydrofluorocarbons, have a more immediate impact on global temperatures than notionally equivalent emissions of  $CO_2$ , and less impact after 20–40 years. Hence, if the primary goal of climate policy is to limit peak warming, then given the time likely to be required to reduce net global  $CO_2$  emissions to zero to stabilize temperatures, the conventional use of  $GWP_{100}$  to compare pulse emissions of  $CO_2$  and SLCPs is likely to overstate the importance of SLCPs for peak warming until global  $CO_2$  emissions are falling<sup>7,8</sup>.

This is not an argument for delay in SLCP mitigation<sup>26</sup>—the benefits to human health and agriculture alone would justify many proposed SLCP mitigation measures<sup>4</sup>—but it is an argument for clarity in what immediate SLCP reductions may achieve for global climate. The use of  $GWP_{100}$  to compare emission pulses might still be appropriate to other policy goals, such as limiting the rate of warming over the coming decades, although the impact of policies on warming rates even over multi-decadal timescales should always be considered in the context of internal climate variability<sup>27</sup>. Some contributions to the rate of sea-level rise also scale with integrated climate forcing<sup>22</sup>.


Simply adopting a different metric that assigns a lower weight to SLCP emissions, such as  $GTP_{100}$ , does not solve this overstatement problem, because any metric that correctly reflects the impact of SLCPs on temperatures 100 years in the future would understate their impact, relative to notionally equivalent quantities of  $CO_2$ , on all shorter timescales. Any choice of metric to compare pulse emissions of cumulative and short-lived pollutants contains a choice

of time horizon<sup>16,18</sup>. It is, however, important for policymakers to be clear about the time horizon they are focusing on. One problem with the  $GWP_{100}$  metric is that ‘warming’ may be interpreted colloquially to mean ‘temperature rise by a point in time’, making the name misleading, because, in the case of SLCPs,  $GWP_{100}$  actually delineates impact on temperatures in 20–40 years, not 100 years.

Figure 2b suggests an alternative way of using  $GWP_{100}$  to express equivalence between cumulative and short-lived climate pollutants that is valid over a wider range of timescales, suggesting a way to use  $GWP_{100}$  to reconcile the ‘emission metrics’ literature<sup>2,3</sup> with the ‘carbon budget’ approach<sup>9</sup>. The solid lines show the impact on global temperatures of a sustained emission of 38 Gt of  $CO_2$ -equivalent ( $CO_2e$ ; again computed using  $GWP_{100}$ ) of the short-lived climate pollutants shown in Fig. 2a, but now starting abruptly in year 1 and distributed evenly over the  $GWP_{100}$  time horizon: hence a sustained emission rate of  $38/(H \times GWP_{100})$  billion tonnes per year, where  $H = 100$  years. These cause temperatures to increase and then approach stabilization after 20–40 years, depending on their lifetimes. The dotted line shows the impact of a pulse emission of 38 Gt $CO_2$  in year one, reproduced from Fig. 2a. The correspondence between these temperature responses is not exact, but much better than in Fig. 2a, at least over timescales from 30 to 100 years.

The reason is simple: a pulse emission of an infinite-lifetime gas and a sudden step change in the sustained rate of emission of a very-short-lifetime gas both give a near-constant radiative forcing. If the total quantities emitted of both gases over the 100-year  $GWP_{100}$  time horizon is the same in terms of  $GWP_{100}$ , then the size of this radiative forcing, and hence the temperature response, will be identical (see Methods for a more formal derivation). The solid and dotted lines in Fig. 2b do not coincide exactly, because  $CO_2$  is not simply an infinite-lifetime gas, nor are the lifetimes of methane or black carbon completely negligible, although the effective residence times of  $CO_2$  and these SLCPs are, crucially, much longer and much shorter, respectively, than the 100-year  $GWP_{100}$  time horizon.

A corollary is that a sustained step change in the rate of emission of a cumulative pollutant such as  $CO_2$  is approximately equivalent to a progressive linear increase or decrease in the rate of emission of an SLCP. This is illustrated in Fig. 2c, which compares the impact of a sustained emission of 38 Gt per year of  $CO_2$  emissions (red dotted line) with SLCP emissions increasing from zero at a rate of 0.38 Gt $CO_2e$  per year per year (solid lines). Again, although the correspondence is not exact, it is much better than the nominally equivalent emission pulses in Fig. 2a. The green dotted line shows that sustained emissions of cumulative pollutants ( $N_2O$  and  $CO_2$ ) have similar impacts on these timescales. Finally, a progressive



**Figure 2 | Impact of pulse versus sustained emissions of various climate forcing agents on global average temperatures.** The colours in each panel indicate the different GHG, with dark grey lines indicating the combined impact of reflective organic and black carbon aerosols (see Methods). **a**, Warming caused by a pulse emission in 2011 with each pulse size being nominally equivalent, using  $\text{GWP}_{100}$ , to 2011 emissions of  $\text{CO}_2$ . **b**, Solid lines show the impact of sustained emissions of SLCPs at a rate equivalent to 2011 emissions of  $\text{CO}_2$  spread over the 100-year  $\text{GWP}_{100}$  time horizon. The dotted red line shows the impact of the pulse emission of  $\text{CO}_2$  reproduced from **a**. **c**, Solid lines show the impact of SLCP emissions progressively increasing from zero at  $0.38 \text{ GtCO}_2\text{e per year per year}$ . Dotted lines show the impact of sustained emissions of  $\text{CO}_2$  (red) and  $\text{N}_2\text{O}$  (green) at  $38 \text{ GtCO}_2$  (or equivalent) per year. **d**, Impact of actual 2011 emissions of each climate forcing agent expressed as a pulse. **e**, Impact of emissions sustained indefinitely at 2011 rates.

change in the rate of emission of  $\text{CO}_2$ , necessary to reach net zero<sup>10</sup>  $\text{CO}_2$  emissions to stabilize temperatures, could only be equated to an accelerating change in SLCP emissions. This last equivalence will eventually become moot because attempting to match the rates of reduction of  $\text{CO}_2$  emissions<sup>28</sup> required to limit warming to  $2^\circ\text{C}$  would result in SLCP emissions having to be reduced below zero. In summary, therefore, a pulse (or sustained) emission of a cumulative pollutant may be approximately equivalent to a sustained (or progressively increasing) change in the rate of emission of an SLCP, but there is no substitute for a progressive reduction in the rate of emission of a cumulative pollutant such as  $\text{CO}_2$ , which remains the *sine qua non* of climate stabilization.

This correspondence between pulse emissions of cumulative pollutants and sustained emissions of short-lived pollutants (or the benefits of corresponding emissions reductions) has been noted before<sup>7,8,11–13</sup>, but previous studies suggested that a new metric of sustained emission reductions would be required to relate them. Figure 2b suggests that the familiar  $\text{GWP}_{100}$  might still be adequate for this purpose, provided it is used to relate sustained reductions in emission rates of SLCPs (agents with lifetimes much shorter than the GWP time horizon) with temporarily avoided emissions of cumulative climate pollutants (any with lifetimes substantially longer than the GWP time horizon).

There are obvious challenges to incorporating this second use of  $\text{GWP}_{100}$  into the UNFCCC process. The Kyoto Protocol and most emissions trading schemes are predicated on emissions accounting over fixed commitment periods. Although possible in the new, more flexible, NDC framework, equating an open-ended commitment to a permanent reduction in an SLCP emission rate with actual avoided emissions of a cumulative pollutant within a commitment period would be a significant policy innovation. Nevertheless, this approximate equivalence may be useful in setting national

or corporate climate policy priorities, particularly where decisions involve capital investments committing future emissions<sup>13</sup>.

This second use of  $\text{GWP}_{100}$  is also relevant to the long-term goal in the Paris Agreement ‘to achieve a balance between anthropogenic emissions by sources and removals by sinks’ to hold the increase in the global average temperature to well below  $2^\circ\text{C}$  above pre-industrial levels. Peak warming scales approximately with cumulative  $\text{CO}_2$  and  $\text{N}_2\text{O}$  emissions (expressed as  $\text{GtCO}_2\text{e}$  using  $\text{GWP}_{100}$ ) between now and the time of peak warming plus the sustained rate of emission of SLCPs (expressed in  $\text{GtCO}_2\text{e}/H$  per year, with  $H = 100$  years if  $\text{GWP}_{100}$  is used to define  $\text{GtCO}_2\text{e}$ ) in the decades immediately before peak warming. So a sustained emission rate of 0.01 tonnes per year of methane has the same impact on peak warming as a pulse of 28 tonnes of  $\text{CO}_2$  released at any time between now and when temperatures peak,  $\text{GWP}_{100}$  of methane being 28. As NDCs are updated, it would be useful for countries to clarify how they propose to balance (individually or collectively) cumulative emissions of  $\text{CO}_2$  and  $\text{N}_2\text{O}$  as these are reduced to zero or below with future emission rates of SLCPs.

Figure 2d shows the impact on global temperatures of actual 2011 emissions of various climate pollutants, considered as a one-year emission pulse<sup>16</sup>. Methane and black carbon emissions in 2011 have a comparable or even larger impact on global temperatures over the next couple of decades than 2011  $\text{CO}_2$  emissions, but their impact decays rapidly, whereas the impact of current  $\text{CO}_2$  emissions persists throughout the twenty-first century and for many centuries beyond.

Figure 2e shows the impact of 2011 emissions of various climate pollutants, assuming these emissions are maintained at the same level for the next 100 years. The warming impact of the cumulative pollutants,  $\text{CO}_2$  and nitrous oxide, increases steadily as long as these emissions persist, whereas sustained emissions of methane and organic and black carbon aerosols cause temperatures to

warm rapidly at first, and then stabilize. A permanent reduction of 50–75% in these SLCPs could reduce global temperatures by over 0.5 °C by mid-century<sup>4</sup>, comparable to the impact on these timescales of similar-magnitude reductions of CO<sub>2</sub> emissions and, it has been argued, at much lower cost<sup>4,5,29</sup>. Stabilizing global temperatures, however, requires net emissions of cumulative pollutants, predominantly CO<sub>2</sub>, to be reduced to zero.

The notion of CO<sub>2</sub>-equivalent pulse emissions of cumulative and short-lived climate pollutants will always be ambiguous because they act to warm the climate system in fundamentally different ways. So far, this ambiguity may have had only a limited impact, not least because emission reductions have so far been relatively unambitious. As countries with relatively large agricultural emissions of methane and significant black carbon emissions begin to quantify their contributions to the UNFCCC, and as the stringency of commitments increases consistent with the collective goal of limiting warming to ‘well below’ 2 °C, this situation may change<sup>21,30</sup>.

For their long-term climate implications to be clear, policies and NDCs need to recognize these differences. GWP<sub>100</sub> can be used in the traditional way, comparing pulse emissions of different GHG, to specify how mitigation of both short-lived and cumulative climate pollutants may reduce the rate and magnitude of climate change over the next 20–40 years, but only over that time. To achieve a balance between sources and sinks of GHG in the very long term, net emissions of cumulative pollutants such as CO<sub>2</sub> need to be reduced to zero, whereas emissions of SLCPs simply need to be stabilized. GWP<sub>100</sub> can again be used, but in the second way identified here, to relate cumulative (positive and negative) emissions of CO<sub>2</sub> until these reach zero with future emission rates of SLCPs, particularly around the time of peak warming. Some NDCs are already providing a breakdown in terms of cumulative and short-lived climate pollutants, or differential policy instruments for different forcing agents<sup>30</sup> and different timescales, all of which is needed for their climatic implications to be clear. The Paris Agreement proposes that Parties will report emissions and removals using common metrics, but a generic CO<sub>2</sub>-equivalent emission reduction target by a given year, defined in terms of GWP<sub>100</sub> and containing a substantial element of SLCP mitigation, represents an ambiguous commitment to future climate. The conventional use of GWP<sub>100</sub> to compare pulse emissions of all gases is an effective metric to limit peak warming if and only if emissions of all climate pollutants, most notably CO<sub>2</sub>, are being reduced such that temperatures are expected to stabilize within the next 20–40 years. This expected time to peak warming will become clear only when CO<sub>2</sub> emissions are falling fast enough to observe the response. Until such a clear end point is in sight, only a permanent change in the rate of emission of an SLCP can be said to have a comparable impact on future temperatures as a one-off pulse emission of CO<sub>2</sub>, N<sub>2</sub>O or other cumulative pollutant.

## Methods

Methods and any associated references are available in the [online version of the paper](#).

Received 26 October 2015; accepted 23 March 2016; published online 2 May 2016

## References

1. *Adoption of the Paris Agreement* FCCC/CP/2015/L.9/Rev.1 (UNFCCC, 2015).
2. Shine, K., Fuglestedt, J., Hailemariam, K. & Stuber, N. Alternatives to the global warming potential for comparing climate impacts of emissions of greenhouse gases. *Climatic Change* **68**, 281–302 (2005).
3. Fuglestedt, J. S. *et al.* Assessment of transport impacts on climate and ozone: metrics. *Atmos. Environ.* **44**, 4648–4677 (2010).
4. Shindell, D. *et al.* Simultaneously mitigating near-term climate change and improving human health and food security. *Science* **335**, 183–189 (2012).

5. Victor, D. G., Kennel, C. F. & Ramanathan, V. The climate threat we can beat. *Foreign Aff.* **91**, 112–114 (2012).
6. Rogelj, J. *et al.* Disentangling the effects of CO<sub>2</sub> and short-lived climate forcer mitigation. *Proc. Natl Acad. Sci. USA* **111**, 16325–16330 (2014).
7. Bowerman, N. H. A. *et al.* The role of short-lived climate pollutants in meeting temperature goals. *Nature Clim. Change* **3**, 1021–1024 (2013).
8. Pierrehumbert, R. T. Short-lived climate pollution. *Annu. Rev. Earth Planet. Sci.* **42**, 341–379 (2014).
9. Allen, M. R. *et al.* Warming caused by cumulative carbon emissions towards the trillionth tonne. *Nature* **458**, 1163–1166 (2009).
10. Matthews, H. D. & Caldeira, K. Stabilizing climate requires near-zero emissions. *Geophys. Res. Lett.* **35**, GL032388 (2008).
11. Smith, S. M. *et al.* Equivalence of greenhouse-gas emissions for peak temperature limits. *Nature Clim. Change* **2**, 535–538 (2012).
12. Lauder, A. R. *et al.* Offsetting methane emissions—an alternative to emission equivalence metrics. *Int. J. Greenh. Gas Contr.* **12**, 419–429 (2013).
13. Alvarez, R. A. *et al.* Greater focus needed on methane leakage from natural gas infrastructure. *Proc. Natl Acad. Sci. USA* **109**, 6435–6440 (2012).
14. *Ecuador's Intended Nationally Determined Contribution* (Gobierno Nacional de la Republica del Ecuador, 2015); <http://www4.unfccc.int/submissions/INDC>
15. Bond, T. C. *et al.* Bounding the role of black carbon in the climate system: a scientific assessment. *J. Geophys. Res.* **118**, 5380–5552 (2013).
16. Myhre, G. *et al.* in *Climate Change 2013: The Physical Science Basis* (eds Stocker, T. F. *et al.*) Ch. 8 (IPCC, Cambridge Univ. Press, 2013).
17. *Brazil's Intended Nationally Determined Contribution* (Federative Republic of Brazil, 2015); <http://www4.unfccc.int/submissions/INDC>
18. Shine, K. P. *et al.* Comparing the climatic effects of emissions of short- and long-lived climate agents. *Phil. Trans. R. Soc. A* **365**, 1903–1914 (2007).
19. Olivie, D. J. L. & Peters, G. P. Variation in emission metrics due to variation in CO<sub>2</sub> and temperature impulse response functions. *Earth Syst. Dyn.* **4**, 267–286 (2013).
20. Reisinger, A. *et al.* Future changes in global warming potentials under representative concentration pathways. *Environ. Res. Lett.* **6**, 024020 (2011).
21. Daniel, J. S. *et al.* Limitations of single-basket trading: lessons from the Montreal Protocol for Climate Policy. *Climatic Change* **111**, 241–248 (2012).
22. Johansson, D. Economics- and physical-based metrics for comparing greenhouse gases. *Climatic Change* **110**, 123–141 (2012).
23. Peters, G., Aamaas, B., Berntsen, T. & Fuglestedt, J. The integrated global temperature change potential (iGTP) and relationships between emission metrics. *Environ. Res. Lett.* **6**, 044021 (2011).
24. Millar, R. J. *et al.* Model structure in observational constraints on the transient climate response. *Climatic Change* **131**, 199–211 (2015).
25. Gillett, N. P. & Matthews, H. D. Accounting for carbon cycle feedbacks in a comparison of the global warming effects of different greenhouse gases. *Environ. Res. Lett.* **5**, 034011 (2010).
26. Schmale, J. *et al.* Air pollution: clean up our skies. *Nature* **515**, 335–337 (2014).
27. Deser, C. *et al.* Uncertainty in climate change projections: the role of internal variability. *Clim. Dynam.* **38**, 527–546 (2010).
28. Allen, M. R. & Stocker, T. F. Impact of delay in reducing carbon dioxide emissions. *Nature Clim. Change* **4**, 23–26 (2013).
29. Huntingford, C. *et al.* The implications of carbon dioxide and methane exchange for the heavy mitigation RCP2.6 scenario under two metrics. *Environ. Sci. Policy* **51**, 77–87 (2015).
30. *New Zealand's Intended Nationally Determined Contribution* (New Zealand Government, 2015); <http://www4.unfccc.int/submissions/INDC>

## Acknowledgements

M.R.A. was supported by the Oxford Martin Programme on Resource Stewardship. M.R.A. and K.P.S. received support from the UK Department of Energy and Climate Change under contract no. TRN/307/11/2011; J.S.F. from the Norwegian Research Council, project no. 235548; R.T.P. from the Kung Carl XVI Gustaf 50-års fond; P.M.F. from the UK Natural Environment Research Council grant no. NE/N006038/1. The authors would like to thank numerous colleagues, particularly among IPCC authors, for discussions of metrics over recent years, and J. Cook for encouraging this work.

## Author contributions

M.R.A. conceived and led the study; all authors contributed to extensive discussions, analysis, interpretation and writing of the paper.

## Additional information

Reprints and permissions information is available online at [www.nature.com/reprints](http://www.nature.com/reprints). Correspondence and requests for materials should be addressed to M.R.A.

## Competing financial interests

The authors declare no competing financial interests.

**Methods**

**Derivation of the approximate equality between GWP<sub>100</sub> and GTP<sub>20-40</sub>.** The equality of GWP<sub>100</sub> and GTP<sub>20-40</sub> follows from the idealized expressions for GWP and GTP for a pulse emission given in ref. 2 (equations A1 and 3 in ref. 2, expressed as relative GWP and GTP respectively, and with decay times replaced by decay rates):

$$GWP_H = \frac{(F_1/k_1)(1 - e^{-k_1 H})}{(F_0/k_0)(1 - e^{-k_0 H})} \tag{1}$$

and

$$GTP_{H'} = \frac{(F_1/(k_1 - k_T))(e^{-k_T H'} - e^{-k_1 H'})}{(F_0/(k_T - k_0))(e^{-k_0 H'} - e^{-k_T H'})} \tag{2}$$

where  $F_1$  is the instantaneous forcing per unit emission and  $k_1$  the concentration decay rate for a GHG, with  $F_0$  and  $k_0$  the corresponding parameters for a reference gas,  $k_T$  is a typical thermal adjustment rate of the ocean mixed layer in response to forcing, and  $H$  and  $H'$  are the GWP and GTP time horizons. For a very short-lived GHG and very long-lived reference gas, such that  $k_1 H \gg 1$ ,  $k_1 H' \gg 1$ ,  $k_0 H \ll 1$ ,  $k_0 H' \ll 1$  and  $k_1 \gg k_T \gg k_0$ , the terms in parentheses in the numerator and denominator of equations (1) and (2) are approximately unity,  $k_0 H$ ,  $e^{-k_T H'}$  and  $(1 - e^{-k_T H'})$  respectively. Hence, using  $k_1 - k_T \approx k_1$  and  $k_T - k_0 \approx k_T$ , we have

$$GWP_H \approx \frac{F_1}{F_0 k_1 H} \quad \text{and} \quad GTP_{H'} \approx \frac{F_1 k_T}{F_0 k_1 (e^{k_T H'} - 1)}$$

so  $GWP_H$  equals  $GTP_{H'}$  if  $H' = \ln(1 + Hk_T)/k_T$ , or 21 years if  $H = 100$  years and  $k_T = (8.4 \text{ years})^{-1}$ , as in ref. 16. Hence, in the limit of a very short-lived gas and infinitely persistent reference gas, the GTP for a pulse emission evaluated at 21 years will be equal to the GWP<sub>100</sub>. The expression becomes more complicated if  $k_1 H' \approx 1$ , as is the case for methane, but this limiting case serves to show that the equality of GWP<sub>100</sub> and GTP<sub>20-40</sub> arises primarily from the thermal adjustment time of the climate system.

**Equivalence between transitory and sustained emissions.** The approximate equivalence of the temperature response to a one-tonne transitory pulse emission of a cumulative pollutant to sustained step change in the rate of emission of an SLCP by  $1/(H \times GWP_H)$  tonnes per year, where  $H$  is the GWP time horizon, follows from the cumulative impact of CO<sub>2</sub> emissions on global temperatures. This means that the temperature response at a time  $H$  after a unit pulse emission of CO<sub>2</sub> (AGTP<sub>p</sub>(CO<sub>2</sub>) in ref. 2), multiplied by  $H$ , is approximately equal to the response after time  $H$  to a one-unit-per-year sustained emission of CO<sub>2</sub> (AGTP<sub>s</sub>(CO<sub>2</sub>)), provided  $H$  is shorter than the effective atmospheric residence time of CO<sub>2</sub>, which is of order millennia. This is consistent with the concept of the ‘trillionth tonne’—that it is the cumulative amount of CO<sub>2</sub> that is emitted, rather than when it is emitted, that matters most for future climate<sup>9</sup>. Ref. 2 also notes that the ratio AGTP<sub>s</sub>( $x$ )/AGTP<sub>s</sub>(CO<sub>2</sub>) is approximately equal to GWP<sub>H</sub>( $x$ ) for time horizons  $H$  much longer than the lifetime of an agent  $x$ . Hence:

$$AGTP_s(x) \approx GWP_H(x) \times AGTP_s(\text{CO}_2) \approx GWP_H(x) \times H \times AGTP_p(\text{CO}_2) \tag{3}$$

provided  $H$  is shorter than the effective residence time of CO<sub>2</sub> and longer than the lifetime of the agent  $x$ , as is the case when  $H = 100$  years and  $x$  is an SLCP.

**The interpretation of an ‘avoided emission pulse.’** Although central to most emission trading schemes, the concept of ‘emission pulse’ may be ambiguous in the context of many mitigation decisions, which may involve policies resulting in permanent changes in emission rates. Another way of expressing this notion of an avoided pulse is in terms of the impact of delay in reducing emissions of cumulative pollutants: a five-year delay in implementing a one-tonne-per-year reduction of CO<sub>2</sub> emissions would need to be compensated for by a permanent reduction of  $5/(100 \times 28) = 1.8 \times 10^{-3}$  tonnes per year of methane (GWP<sub>100</sub> of methane being 28). This would compensate only for the direct impact of the delay in CO<sub>2</sub> emission reductions, not for additional committed future CO<sub>2</sub> emissions that might also result from that delay<sup>28</sup>.

**Treatment of black carbon emissions.** Focusing solely on absorbing aerosols gives a high estimated ‘radiative efficiency’ (impact on the global energy budget per unit change in atmospheric concentration) for black carbon, a strong positive global climate forcing<sup>15</sup> (1.1 W m<sup>-2</sup> in 2011) and a GWP<sub>100</sub> of 910. This figure has been argued<sup>16</sup> to be too high, and the actual radiative impact of individual black carbon emissions depends strongly on the circumstances (location, season and weather conditions) at the time of emission. Many processes that generate black carbon also generate reflective organic aerosols, which have a cooling effect on global climate. Although ratios vary considerably across sources, policy interventions to limit black carbon emissions are likely also to affect these other aerosols, so it might be more relevant to consider their combined impact: the current best estimate<sup>16</sup> net global radiative forcing of organic and black carbon aerosols in 2011 was 0.35 W m<sup>-2</sup>, giving a combined GWP<sub>100</sub> of 290, used in the figures. Combined emissions of organic and black carbon aerosols are inferred from this GWP<sub>100</sub> value, assuming all radiative forcing resulting from these emissions is concentrated in the first year (that is, a lifetime much shorter than one year). This is only one estimate of a very uncertain quantity: when both reflection and absorption are taken into account, including interactions between aerosols and clouds and surface albedo, even the sign of the net radiative impact of the processes that generate black carbon aerosols remains uncertain.

**Modelling details.** For Fig. 1 GWP values were calculated using current IPCC methane and CO<sub>2</sub> impulse response functions without explicit carbon-cycle feedbacks<sup>16</sup>. Radiative forcing (RF) of a pulse emission of organic and black carbon aerosols concentrated in year 1, scaled to give a net GWP<sub>100</sub> of 290, consistent with ratio of 2011 RF values given in refs 15,16. GTP values were calculated using the standard IPCC AR5 thermal response model (solid blue lines) with coefficients adjusted (dotted blue lines) to give realized warming fractions<sup>24</sup> (ratio of the transient climate response (TCR) to equilibrium climate sensitivity (ECS)) of 0.35 and 0.85, spanning the range of uncertainty around the best-estimate value of 0.56. Values in Fig. 2 were calculated as for Fig. 1 with radiative efficiencies and lifetimes provided in Table A.8.1 of ref. 16 and representative mid-range values of TCR = 1.5 °C and ECS = 2.7 °C.